

Policy Brief: Climate change impacts of plastics

Nihan Karali, Ellen Palm, Juan Baztan, Patricia Villarubia Gomes, Nina Khanna, Karin Kvale, Ana Luzia Lacerda, Bethany Jorgenson

▶ To cite this version:

Nihan Karali, Ellen Palm, Juan Baztan, Patricia Villarubia Gomes, Nina Khanna, et al.. Policy Brief: Climate change impacts of plastics. 2024. hal-04423053

HAL Id: hal-04423053 https://uvsq.hal.science/hal-04423053v1

Preprint submitted on 29 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Policy Brief:

Climate change impacts of plastics

The environmental impact of plastics is multi-fold, contributing to the triple planetary crisis of climate change, biodiversity loss, and pollution. Plastic pollution and climate change are intrinsically interconnected issues. The plastics value chain has a significant climate impact throughout the life cycle from extraction of fossil fuel to production to end-of-life disposal. However, up- and downstream greenhouse gas (GHG) emissions from plastics production and disposal are often not taken into consideration and may have been overlooked in the INC negotiation process. This Policy brief delves into the ways in which plastic contributes to the emission of GHGs throughout its life cycle.

Climate change impact of the plastics life cycle

Studies estimate that plastics account for around 3-8% of global GHG emissions, and it is projected to double by 2060 [1-4].

- Most of the GHG emissions from the plastic industry are in the form of carbon dioxide (CO₂) and are related to the energy used at various stages of the plastic life cycle, including extraction of fossil fuels, production of monomers, resins, and additives, plastic conversion, and end-of-life management. There are also process emissions that include CO₂ as well as other GHGs (e.g., fugitive methane emissions from upstream oil and gas operations) [1,5].
- 99% of plastic polymers and chemical additives are made from fossil fuel feedstocks [6]. Almost 15% of crude oil production goes to the petrochemicals sector with plastics as the largest end-use product [7]. Combustion of oil and gas is also required for power and high-temperature heat demand in the production of plastics.
- Estimates indicate that plastics and other petrochemicals industries will drive half of the growth in demand of fossil fuel production by 2050 [7]. The extraction, refining, and transportation of petrochemicals for plastic production involves various energy and emission-intensive processes.
- However, most studies of resin or monomer production ignore previous steps, thus providing a rather incomplete picture and underestimating the full climate impact [8].
- Studies estimate that emissions from end-of-life processes contribute to around 10% of total emissions from the plastic lifecycle, mainly from incineration. While production is the major concern for climate impact, continued mismanagement of end-of-life plastics presents a significant GHG emissions issue going forward [1,9].
- Incineration, in which about 20% of the plastic waste currently ends, is an energy and emission-intensive process [4]. Waste-to-energy is another form of incineration that uses plastic waste as an energy source [10]. However, burning fossil fuel-based plastic waste is not better for the climate than burning other fossil fuels [2,9]. It also generates toxic air pollution that is harmful to human health due to the chemicals released, and is particularly harmful to often-adjacent lower-income communities already impacted by climate change [11]. Some of these emissions could possibly be captured with carbon capture technologies, but these technologies remain largely unproven at scale, with concerns of cost and long-term mitigation impact [12,13].

- Mechanical recycling can lead to GHG emission reductions by avoiding incineration and reducing
 virgin plastic production. However, the quality of the material is often reduced, so the final product
 may not provide the same functionality [14]. Because of degradation, mechanically recycled products
 may still end up being incinerated after a few cycles, which must be avoided through other end-oflife treatment processes [15]. Delayed incineration negates short-term emission reductions and is
 not part of a sustainable recycling system [9,15]
- Chemical recycling processes are energy and emission-intensive [16,17], and depending on the
 recycling process (e.g. feedstock recycling, depolymerization and purification), the emissions
 intensity of recycling may be higher than that of virgin plastic production. Chemical recycling of
 plastics is also linked to generation of toxic materials [11].
- Landfill waste, in which about 40% of plastic waste current ends, is another source of GHG emissions with potential for additional GHG leakage with decomposition over time [4,18].

How can the treaty include climate change?

The combined effects of plastics pollution and climate change can have a disproportionate impact on already vulnerable populations through increased exposure to pollution, extreme weather events, and the health risks associated with the spread of vector diseases and heat exposure [19].

To simultaneously reduce plastic pollution and the climate change impacts of plastics the plastic treaty could consider designing strategies, such as control measures across the lifecycle of plastic, that weigh the overall impacts of alternate upstream and downstream trajectories, including but not limited to plastic production caps, reduce, reuse, and recycling.

Current plastic production is contributing to the devastating human health consequences associated with the triple planetary crisis of climate change, biodiversity loss, and pollution [4,20]. A robust and science-based approach is essential if the plastics treaty is to be effective, providing a narrative rooted in the scientific evidence explicitly addressing the consequences of fossil fuel extraction, conversion, and disposal, and offering transparent guidance on potential pathways for global cooperation and action towards a more sustainable and healthy future.

Contributors and references

This briefing was prepared by members of the Scientists' Coalition for an Effective Plastics Treaty:

Authors: Nihan Karali (co-chair) (Lawrence Berkeley National Laboratory, US), Ellen Palm (co-chair) (Lund University, Sweden), Juan Baztan (CEARC, Université de Versailles SQY, France), Patricia Villarrubia Gomez (Stockholm Resilience Centre, Stockholm University, Sweden), Nina Khanna (Lawrence Berkeley National Laboratory, US), Karin Kvale (GNS Science, New Zealand), Ana Luzia Lacerda (Sorbonne University, France), Bethany Jorgensen (Cornell University, US)

Reviewers: Fredric Bauer, Kishore Boodhoo, Xuejing Chen, Grau Etienne, Annika Jahnke, Jacob Kean-Hammerson, Kristoffer Kortsen, Doris Knoblauch, Eva Kumar, Peter Stoett, Hideshige Takada, Melissa Wang, Roland Weber.

Contact: scientists.coalition@ikhapp.org

References

- [1] Zheng, J. and Suh, S. (2019) Strategies to reduce the global carbon footprint of plastics. Nat. Clim. Chang. 9, 374–37.
- [2] Meng et al. (2023) Planet-compatible pathways for transitioning the chemical industry. PNAS. 120 (8) e2218294120.
- [3] CIEL (2019) Plastic & climate: the hidden costs of a plastic planet.

- [4] OECD (2022) Global Plastics Outlook Policy Scenarios to 2060.
- [5] IPCC (2022) Climate Change 2022 Mitigation of Climate Change Working Group III Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change Summary for Policymakers.
- [6] Levi and Cullen (2018) Mapping Global Flows of Chemicals: From Fossil Fuel Feedstocks to Chemical Products. Environmental Science & Technology 2018 52 (4), 1725-1734.
- [7] IEA (2018), The Future of Petrochemicals, IEA, Paris. License: CC BY 4.0.
- [8] EIA (2022) Convention on Plastic Pollution Essential Elements: Virgin Plastic Production and Consumption.
- [9] Stegmann et al. Plastic futures and their CO2 emissions. Nature 612, 272–276 (2022).
- [10] Vlasopoulos et al. (2023) Life cycle assessment of plastic waste and energy recovery. Energy, Volume 277, 127576.
- [11] UNEP (2023) Chemicals in Plastics A Technical Report.
- [12] Istrate et al. (2023) Prospective analysis of the optimal capacity, economics and carbon footprint of energy recovery from municipal solid waste incineration. Resources, Conservation and Recycling. Volume 193, 106943.
- [13] Kleijne et al. (2022) Limits to Paris compatibility of CO2 capture and utilization. One Earth. Volume 5 (2), 168-185.
- [14] Schyns and Shaver (2020) Mechanical Recycling of Packaging Plastics: A Review. Macromolecular Rapid Communications Volume 42, Issue 3: Polymers for a Sustainable Future.
- [15] Kortsen et al. (2023) A plastics hierarchy of fates: sustainable choices for a circular future. arxiv.org/abs/2303.14664
- [16] Material Economics (2019). Industrial Transformation 2050 Pathways to Net-Zero Emissions from EU Heavy Industry.
- [17] Meys et al. (2020) Towards a circular economy for plastic packaging wastes the environmental potential of chemical recycling. Resources, Conservation and Recycling. Volume 162, 105010.
- [18] Tenhunen-Lunkka et al. (2023) Greenhouse Gas Emission Reduction Potential of European Union's Circularity Related Targets for Plastics. Circ.Econ.Sust. 3, 475–510.
- [19] The Minderoo-Monaco Commission on Plastics and Human Health (2023) Annals of Global Health. 21: 89 (1).
- [20] Bauer et al. (2022) Plastics and climate change—Breaking carbon lock-ins through three mitigation pathways. One Earth, Volume 5, Issue 4.

Please cite this as: Scientists' Coalition for an Effective Plastics Treaty (2023) Policy Brief: Climate change impacts of plastics. DOI: 10.5281/zenodo.7972056