

Data-driving methods: More than merely trendy buzzwords?

Julien Textoris, Fabio Silvio Taccone, Lara Zafrani, Antoine Guillon, Sébastien Gibot, Fabrice Uhel, Eric Azabou, Guillaume Monneret, Frédéric Pène, Nicolas de Prost, et al.

▶ To cite this version:

Julien Textoris, Fabio Silvio Taccone, Lara Zafrani, Antoine Guillon, Sébastien Gibot, et al.. Data-driving methods: More than merely trendy buzzwords?. Annals of Intensive Care, 2018, 8 (1), pp.58. 10.1186/s13613-018-0405-7. hal-04532120

HAL Id: hal-04532120 https://uvsq.hal.science/hal-04532120v1

Submitted on 4 Apr 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

REVIEW

Open Access

CrossMark

Data-driving methods: More than merely trendy buzzwords?

Julien Textoris¹, Fabio Silvio Taccone², Lara Zafrani³, Antoine Guillon⁴, Sébastien Gibot⁵, Fabrice Uhel⁶, Eric Azabou⁷, Guillaume Monneret⁸, Frédéric Pène⁹, Nicolas de Prost^{10*} and Stein Silva¹¹ on behalf of the Translational Research Committee of the French Intensive Care Society (Société de Réanimation de Langue Française, SRLF)

Keywords: Big data, Machine learning, Artificial intelligence, Critical care

Intensive care units (ICU) physicians are experiencing a rapidly expanding collection of vast amounts of data from routine practice, patients' monitoring as well as from diagnostic or prognostic tests. However, although these data could influence their clinical decisions and management, the validity and relevance of data processing methods, in particular in case of complex data sets (i.e. so-called big data, see Table 1 for related terminology) remain to be defined. A growing body of research has recently suggested that emerging artificial intelligence (AI)-derived methods could help physicians to access, organize and use important amounts of data more easily. Nowadays, such methods have already found applications in various fields, including technology, biology, computer science or sociology [1]. However, are these approaches more than merely trendy buzzwords? Are they reliable enough to match the exponential growth of medical complexity in the critical care setting? And, last but not least, can the holistic use of massive data sources available eventually provide clinically relevant information?

The reality is that the exponential combinations of patients, conditions and treatments cannot be exhaustively explored by processes that often—intentionally or inadvertently—exclude interdependent input/output parameters because they do not fit into a priori hypotheses or predefined models (Additional file 1: Figure S1). In such a context, data-driven approaches hold promise of accurately dealing with big data methodological issues, and doing so might have a significant impact on the improvement in diagnosis, monitoring and prognostication of ICU processes.

ICU database: closing the data loop

As an evolution to this approach, a dynamic clinical "data mining" (Table 1) has been recently proposed, based on "data-driven" methods (Additional file 1: Figure S1). The main idea is the use of feedback loops to enable real-time analysis of patient databases, allow the optimization of patient's care and lead to more efficient targeting of tests, treatments and vigilance for adverse effects (e.g. "Multiparameter Intelligent Monitoring in Intensive Care" (MIMIC) [2]. Such closed-loop databases provide physicians with a unique opportunity to accumulate useful clinical evidence to: (1) identify patient subpopulations with important variations in treatment efficacy or unexpected delayed adverse effects, (2) reveal interactions between simultaneous treatments and physiological conditions, (3) create and cross-validate (Table 1) predictive models across research teams and institutions to better determine which findings are generalizable and (4) pave the way for the development and validation of innovative and more personalized treatments.

Establishing knowledge

Big data methods seem to have straightforward applications for personalized medicine [3] and might pave the way for promising studies focused on the analysis of the intrinsic complexity underpinning human physiology.

¹⁰ Service de Réanimation Médicale, Hôpital Henri Mondor, 51, Avenue du Maréchal de Lattre de Tassigny, 94010 Créteil Cedex, France

Full list of author information is available at the end of the article



[©] The Author(s) 2018. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

^{*}Correspondence: nicolas.de-prost@aphp.fr

Table 1 Data-driven analysis and related terminology


Big data	Data sets with size/complexity beyond the capacity of commonly used methodological approaches to capture, manage and process data. Big data might be defined by their high <i>volume</i> , large <i>variety</i> and the important <i>velocity</i> that is required to process (3v definition)
Closed-loop system	System in which some or all its outputs are used as inputs. In health care, the use of such feedback loop enables real-time analysis of patient databases and could permit to optimize clinical care leading to more efficient targeting of tests and treatments and vigilance for adverse effects (i.e. dynamic clinical data mining)
Cross-validation	Statistical technique for assessing how the results of an analysis will generalize to an independent data set. For example, doing so it could permit to estimate how accurately a predictive model will perform in practice
Crowdsourcing	The practice of obtaining needed solution by soliciting contribution from a large group of people and specially from online com- munities
Data mining	The process of collecting, searching through and analysing a large amount of data in a database, as to discover patterns of rela- tionships. It is worth noting that this approach does not look for causality and simply aim to detect significant data configura- tions
Machine learning	Derived methods from artificial intelligence that provides computers with the ability to learn without being explicitly pro- grammed. The process of machine learning uses the data to detect patterns and adjust programme actions accordingly

Omics: the rise of the narciss-ome

Omics data represent a massive source of multimodal data. The European Bioinformatics Institute (EBI), one of the world's largest biological data repositories, is currently storing: ~5 petabytes (Additional file 2: Table S1) nucleotide sequence data, more than 30,000 genomes and ~2 million gene expression assays [3]. Furthermore, this infrastructure has been accessed 562 million times each month by ~9 million distinct hosts in 2015. These impressive figures highlight the fact that data-driven analysis methods are already a constituent part of worldwide collaborative research projects, built on large big data sharing (Fig. 1a).

Brain, consciousness and complexity

To illustrate this point and demonstrate how data-driven approaches could be successfully used in this setting, we can describe a recent study focused on the assessment of brain structural impact of anoxic/hypoxic insult related to cardiac arrest (CA), and the potential use of brain MRI grey matter morphometry to predict patients' one-year neurological outcome. The authors [4] studied a large and multicenter cohort of anoxic comatose patients, which were scanned during the acute phase following CA in standardized conditions. Crucially, to accurately evaluate whole-brain grey matter morphometry in this setting, fine-grained quantification techniques were applied. Eventually, a data-driven approach was used and



permitted to obtain a predictive classifier that showed a significant discriminative power [4] and enabled the identification of brain grey structures whose degree of atrophy was significantly related to one-year neurological outcome (Fig. 1b).

Promises, pitfalls and challenges

Complex statistical analyses designed to deal with large data sets might appear as magic bullets rendering cumbersome randomized trials dispensable (Additional file 3: Table S2). In fact, we should certainly keep in mind that these statistical optimization techniques are not shortcuts to broader medical reasoning and should not deter clinicians from carefully scrutinizing data so that to avoid inappropriate and naive use of these elegant analytical methods. For example, population selection and adjustment processes may dramatically influence the outcome of studies, giving rise to diametrically opposite conclusions [5].

Furthermore, few additional and unavoidable challenges, which are specifically related to the use of datadriven methods should be addressed: (1) computational issues should be adequately addressed probably by means of cloud storage and cloud computing facilities [6], (2) improving quality and ability to structure data, to ensure interoperability between various sources of data [7], (3) cultural and ethical issues should also be considered and constitute a still moot issue in the field, raising questions on data ownership, patient anonymity, agreement to participate and accountability [8], and highlight the need for further debate, standardization and update of the current legal or regulatory frameworks [9] and (4) finally, it is worth noting that the need for specific analytical skills (inference, prediction and computational abilities) justifies new collaborative interactions between research teams as well as specific training for both data scientist and future physicians [10].

Conclusion

Considering the complexity of ICU setting, we have illustrated how data-driven approaches, through closed-loop systems integrating multimodal data, hold the promise to provide individually tailored and real-time patient care based on the large amount of information currently at our disposal. Regarding translational research, datadriven and hypothesis-driven approaches appear not to be mutually exclusive, but largely complementary and reciprocally challenging. Understanding the opportunities and pitfalls of implementing big data in the ICU setting and considering the subsequent technical, ethical and societal changes are key issues for the upcoming years, paving the way for critical diagnostic and therapeutic innovations.

Additional files

Additional file 1: Figure S1. Analytical methods for biomedical research. Compared to rational hypothesis-driven research methods (upper panel), data-driven analysis (lower panel) does not imply reductions neither of the number of hypothesis that could be studied (i.e. including dynamical interactions), nor of the obtained data that is used to extract relevant information. Additionally, hypothesis-driven methods are built on optimised models derived from artificial intelligence domains, which can learn and evolve without explicit programming, and validate the created model using data from multiple and independent data sets (i.e. machine learning, supplementary-table-1 for related terminology).

Additional file 2: Table S1 Conventional terms used to describe data size. Scale is based on powers of 1000.

Additional file 3: Table S2 Opportunities and difficulties related to datadriven analysis.

Abbreviations

Al: artificial intelligence; CA: cardiac arrest; EBI: European Bioinformatics Institute: ICU: intensive care unit: MIMIC: Multiparameter Intelligent Monitoring in Intensive Care

Authors' contributions

JT and SS drafted the manuscript; all authors read, corrected and approved the final version of the manuscript.

Author details

Département d'Anesthésie-Réanimation, hôpital Édouard-Herriot, Hospices Civils de Lyon, CHU de Lyon, 69437 Lyon, France.² Service de Soins Intensifs, Hôpital Erasme, 1070 Brussels, Belgium.³ Service de Réanimation Médicale, APHP Hôpital Saint-Louis, Paris, France.⁴ Service de Médecine Intensive Réanimation, CHU de Tours, 37000 Tours, France.⁵ Service de Réanimation Médicale, Hôpital Central, CHU de Nancy, 54000 Nancy, France.⁶ Service de Réanimation Médicale et Maladies Infectieuses, Hôpital Pontchaillou, CHU de Rennes, Rennes, France.⁷ Service de Réanimation, APHP Hôpital Raymond Poincaré, Garches, 92380 Paris, France.⁸ Laboratoire d'immunologie, hôpital Edouard Herriot, Hospices Civils de Lyon, CHU de Lyon, 69437 Lyon, France. ⁹ Service de Réanimation Médicale, APHP, Hôpital Cochin, Paris, France.¹⁰ Service de Réanimation Médicale, Hôpital Henri Mondor, 51, Avenue du Maréchal de Lattre de Tassigny, 94010 Créteil Cedex, France.¹¹ Service de Réanimation, CHU Purpan, 31300 Toulouse, France.

Acknowledgements

None

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials

Not applicable

Consent for publication

Not applicable.

Ethics approval and consent to participate Not applicable

Funding

None.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 19 March 2018 Accepted: 23 April 2018 Published online: 02 May 2018

References

- 1. Khanna S, Sattar A, Hansen D. Artificial intelligence in health—the three big challenges. Australas Med J. 2013;6(5):315–7.
- Saeed M, Villarroel M, Reisner AT, Clifford G, Lehman L-W, Moody G, et al. Multiparameter Intelligent Monitoring in Intensive Care II: a public-access intensive care unit database. Crit Care Med. 2011;39(5):952–60.
- Chen R, Mias GI, Li-Pook-Than J, Jiang L, Lam HYK, Chen R, et al. Personal omics profiling reveals dynamic molecular and medical phenotypes. Cell. 2012;148(6):1293–307.
- Silva S, Peran P, Kerhuel L, Malagurski B, Chauveau N, Bataille B, et al. Brain gray matter mri morphometry for neuroprognostication after cardiac arrest. Crit Care Med. 2017;45(8):e763–71.
- Gershengorn HB, Wunsch H, Scales DC, Zarychanski R, Rubenfeld G, Garland A. Association between arterial catheter use and hospital mortality in intensive care units. JAMA Intern Med. 2014;174(11):1746–54.

- 6. Chen J, Qian F, Yan W, Shen B. Translational biomedical informatics in the cloud: present and future. Biomed Res Int. 2013;2013:658925.
- Kruse CS, Goswamy R, Raval Y, Marawi S. Challenges and opportunities of big data in health care: a systematic review. JMIR Med Inform. 2016;4(4):e38.
- Zook M, Barocas S, Boyd D, Crawford K, Keller E, Gangadharan SP, et al. Ten simple rules for responsible big data research. PLoS Comput Biol. 2017;13(3):e1005399.
- 9. Sivarajah U, Kamal MM, Irani Z, Weerakkody V. Critical analysis of big data challenges and analytical methods. J Bus Res. 2017;70:263–86.
- Sejdić E. Education: gear students up for big medical data. Nature. 2015;518(7540):483.

Submit your manuscript to a SpringerOpen[™] journal and benefit from:

- Convenient online submission
- ► Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at > springeropen.com